

Rare B_s decays to η and η' final states

M. V. Carlucci^{a,b}, P. Colangelo^a, F. De Fazio^a

^a *Istituto Nazionale di Fisica Nucleare, Sezione di Bari, Italy*

^b *Dipartimento di Fisica, Università di Bari, Italy*

We study exclusive B_s decays to final states with η and η' , induced by the rare $b \rightarrow s\ell^+\ell^-$ and $b \rightarrow s\nu\bar{\nu}$ transitions. Differential decay rates and total branching fractions are predicted in the Standard Model, adopting the flavour scheme for the description of the η - η' mixing. We discuss the theoretical uncertainty related to the hadronic matrix elements. We also consider these decay modes in a new Physics scenario with a single universal extra dimension, studying the dependence of branching ratios and decay distributions on the compactification scale R^{-1} of the extra dimension.

PACS numbers: 12.60.-i, 13.25.Hw

I. INTRODUCTION

The study of B_s properties and decays plays an important role in the exploration of the Standard Model (SM) and in the searches for new Physics phenomena, and a great experimental effort is devoted at present and foreseen in the near future at the hadron colliders and at the B factories running at the $\Upsilon(5S)$ peak [1]. In particular, $B_s^0 - \bar{B}_s^0$ oscillations provide complementary information with respect to K^0 and B_d systems for the analysis of CP violation. CDF and D0 Collaboration at the Fermilab TeVatron have recently carried out measurements of the $B_s^0 - \bar{B}_s^0$ mixing phase, by an angular analysis of the final state $J/\psi\phi$. The measured phase is larger than the SM prediction (although with a sizeable error) [2]. If confirmed, this result would represent an evidence of Physics beyond SM. For this reason, new measurements are foreseen, involving also the final states $J/\psi f_0(980)$, $J/\psi\eta$ and $J/\psi\eta'$; the branching ratio of $B_s \rightarrow J/\psi\eta$ has been recently measured by the Belle Collaboration using the $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^0\pi^-$ modes to reconstruct η mesons [3].

B_s is of prime interest also for several rare decay modes, namely those induced by the $b \rightarrow s$ transition, that are potentially important for detecting new Physics effects. Here we focus on the decays into η and η' and a pair of leptons, either $\ell^+\ell^-$ (with $\ell = e, \mu$ and τ) or $\nu\bar{\nu}$. Our aim is to give predictions for several observables in the Standard Model, discussing the theoretical uncertainties related to the $\eta - \eta'$ mixing and to the hadronic matrix elements in the decay amplitudes.

A second purpose is to consider a specific new Physics scenario and study how the various observables deviate from SM. The chosen framework is the Appelquist-Cheng-Dobrescu (ACD) model with a single universal extra dimension (UED) [4]. The model is a minimal extension of SM in $4+1$ dimensions, with the extra dimension compactified to the orbifold S^1/Z_2 and the fifth coordinate y running from 0 to $2\pi R$, $y = 0$ and $y = \pi R$

being fixed points of the orbifold. The fields are allowed to propagate in all $4+1$ dimensions, hence the model belongs to the class of *universal* extra dimension scenarios. One of its motivations is the possibility of naturally providing candidates for the dark matter, an issue of fundamental importance.

In the ACD model the SM particles correspond to the zero modes of fields propagating in the compactified extra dimension. In addition to the zero modes, towers of Kaluza-Klein (KK) excitations are predicted to exist, corresponding to the higher modes of the fields in the extra dimension; such fields are imposed to be even under a parity transformation in the fifth coordinate $P_5 : y \rightarrow -y$. On the other hand, fields which are odd under P_5 propagate in the extra dimension without zero modes, and correspond to particles without SM partners.

The masses of KK particles depend on the radius R of the compactified extra dimension, the new parameter with respect to SM [42]. For example, the masses of the KK bosonic modes are given by:

$$m_n^2 = m_0^2 + \frac{n^2}{R^2} \quad n = 1, 2, \dots \quad (1.1)$$

m_0 being the mass of the zero mode, so that for small values of R , i.e. at large compactification scales, the KK particles decouple from the low energy regime. Another property of the ACD model is the conservation of the KK parity $(-1)^j$, j being the KK number. KK parity conservation implies the absence of tree level contributions of Kaluza Klein states to processes taking place at low energy, $\mu \ll 1/R$, forbidding the production of a single KK particle off the interaction of standard particles. This permits to use precise electroweak measurements to provide a lower bound to the compactification scale: $1/R \geq 250 - 300$ GeV [5]. Moreover, this suggests the possibility that the lightest KK particles are among the dark matter components, namely the $n = 1$ Kaluza-Klein excitations of the photon and neutrinos [6, 7].

Since KK modes can affect the loop-induced processes, Flavour Changing Neutral Current (FCNC) transitions are particularly suitable for constraining this new Physics scenario, and indeed many observables are sensitive to the compactification radius in case, e.g., of processes with B and Λ_b [8, 9, 10, 11, 12]. Here we consider rare B_s decays into η and η' mesons: $B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-$ and $B_s \rightarrow \eta^{(\prime)} \nu \bar{\nu}$, described by the effective Hamiltonian reported in Section II. We discuss in Section III the role of the η - η' mixing and of the $B_s \rightarrow \eta^{(\prime)}$ form factors, and present predictions for the various modes in the following two Sections, with comments on the feasibility of measuring properties of these processes. Before concluding, we discuss in the ACD model the $1/R$ dependence of the bs Unitarity Triangle, i.e. the condition among the Cabibbo-Kobayashi-Maskawa (CKM) elements involved in B_s decays, and in particular the possible value of the CP violating phase β_s in this model.

II. EFFECTIVE HAMILTONIAN FOR $B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-$ AND $B_s \rightarrow \eta^{(\prime)} \nu \bar{\nu}$

In the Standard Model, the effective $\Delta B = -1$, $\Delta S = 1$ Hamiltonian describing the transition $b \rightarrow s \ell^+ \ell^-$ can be expressed in terms of a set of local operators:

$$H_{b \rightarrow s \ell^+ \ell^-} = -4 \frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_{i=1}^{10} C_i(\mu) O_i(\mu) \quad (2.1)$$

where G_F is the Fermi constant and V_{ij} are elements of the CKM mixing matrix (terms proportional to $V_{ub} V_{us}^*$ are neglected since the ratio $\left| \frac{V_{ub} V_{us}^*}{V_{tb} V_{ts}^*} \right|$ is $\mathcal{O}(10^{-2})$). The operators O_i are written in terms of quark and gluon fields:

$$\begin{aligned} O_1 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) (\bar{\ell}_{L\beta} \gamma_\mu \ell_{L\beta}) \\ O_2 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\beta}) (\bar{\ell}_{L\beta} \gamma_\mu \ell_{L\alpha}) \\ O_3 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) [(\bar{u}_{L\beta} \gamma_\mu u_{L\beta}) + \dots + (\bar{b}_{L\beta} \gamma_\mu b_{L\beta})] \\ O_4 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\beta}) [(\bar{u}_{L\beta} \gamma_\mu u_{L\alpha}) + \dots + (\bar{b}_{L\beta} \gamma_\mu b_{L\alpha})] \\ O_5 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) [(\bar{u}_{R\beta} \gamma_\mu u_{R\beta}) + \dots + (\bar{b}_{R\beta} \gamma_\mu b_{R\beta})] \\ O_6 &= (\bar{s}_{L\alpha} \gamma^\mu b_{L\beta}) [(\bar{u}_{R\beta} \gamma_\mu u_{R\alpha}) + \dots + (\bar{b}_{R\beta} \gamma_\mu b_{R\alpha})] \\ O_7 &= \frac{e}{16\pi^2} m_b (\bar{s}_{L\alpha} \sigma^{\mu\nu} b_{R\alpha}) F_{\mu\nu} \\ O_8 &= \frac{g_s}{16\pi^2} m_b \left[\bar{s}_{L\alpha} \sigma^{\mu\nu} \left(\frac{\lambda^a}{2} \right)_{\alpha\beta} b_{R\beta} \right] G_{\mu\nu}^a \\ O_9 &= \frac{e^2}{16\pi^2} (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) \bar{\ell} \gamma_\mu \ell \\ O_{10} &= \frac{e^2}{16\pi^2} (\bar{s}_{L\alpha} \gamma^\mu b_{L\alpha}) \bar{\ell} \gamma_\mu \gamma_5 \ell \end{aligned} \quad (2.2)$$

with α, β colour indices, $b_{R,L} = \frac{1 \pm \gamma_5}{2} b$, and $\sigma^{\mu\nu} = \frac{i}{2} [\gamma^\mu, \gamma^\nu]$; e and g_s are the electromagnetic and the strong coupling constant, respectively, and $F_{\mu\nu}$ and $G_{\mu\nu}^a$ in O_7

and O_8 denote the electromagnetic and the gluonic field strength tensor. O_1 and O_2 are current-current operators, $O_3, \dots, O_6$ QCD penguin operators, O_7 and O_8 magnetic penguin operators, O_9 and O_{10} semileptonic electroweak penguin operators. The Wilson coefficients in (2.1) have been computed at NNLO in the Standard Model [13]. The operators O_1 and O_2 contribute to the final state with a lepton pair through a $\bar{c}c$ contribution that can give rise to charmonium resonances J/ψ , $\psi(2S)$, etc. The resonant term can be controlled and subtracted by appropriate kinematical cuts around the resonance masses. Since the Wilson coefficients $C_3 - C_6$ are small, the contribution of only the operators O_7, O_9 and O_{10} can be kept for the description of the $b \rightarrow s \ell^+ \ell^-$ transition, with a modification of the Wilson coefficient C_7 described below.

The SM effective Hamiltonian for $b \rightarrow s \nu \bar{\nu}$:

$$H_{b \rightarrow s \nu \bar{\nu}} = \frac{G_F}{\sqrt{2}} \frac{\alpha}{2\pi \sin^2(\theta_W)} V_{tb} V_{ts}^* \eta_X X(x_t) O_L = C_L O_L \quad (2.3)$$

involves the operator

$$O_L = \bar{s} \gamma^\mu (1 - \gamma_5) b \bar{\nu} \gamma_\mu (1 - \gamma_5) \nu. \quad (2.4)$$

θ_W is the Weinberg angle; the function $X(x_t)$ ($x_t = \frac{m_t^2}{M_W^2}$, with m_t the top quark mass) has been computed in [14] and [15, 16], while the QCD factor η_X is close to one [15, 16, 17], so that we use $\eta_X = 1$.

In the ACD model no operators other than those in (2.1), (2.2) and (2.3) contribute to $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$. The model belongs to the class of Minimal Flavour Violating (MFV) models, and the effects beyond SM are only encoded in the Wilson coefficients of the effective Hamiltonian [8]. KK excitations modify the coefficients C_i and C_L , which acquire a dependence on the compactification scale $1/R$. For large values of $1/R$, due to decoupling of massive KK states, the coefficients C_i and C_L reproduce the Standard Model values and the SM phenomenology is recovered.

The Wilson coefficients can be expressed as functions $F(x_t, 1/R)$ generalizing the SM analogues $F_0(x_t)$:

$$F(x_t, 1/R) = F_0(x_t) + \sum_{n=1}^{\infty} F_n(x_t, x_n), \quad (2.5)$$

with $x_n = \frac{n^2}{R^2 M_W^2}$. Remarkably, the sum over the KK contributions in (2.5) is finite, a consequence of a generalized GIM mechanism [8]; the SM results are recovered for $R \rightarrow 0$, since $F(x_t, 1/R) \rightarrow F_0(x_t)$ in that limit.

For $1/R$ of the order of a few hundreds of GeV the coefficients differ from the Standard Model values, and the physical observables are predicted to be different than in SM. For the exclusive decays, however, it is important to study if this effect can be picked up, or it is obscured by other uncertainties, in particular those affecting the B_s and η, η' matrix elements of the operators in (2.1), (2.3).

III. $\eta - \eta'$ MIXING AND $B_s \rightarrow \eta^{(\prime)}$ FORM FACTORS

In the determination of the $B_s \rightarrow \eta^{(\prime)}$ matrix elements we need to account for the $\eta - \eta'$ mixing. This is usually described in two schemes: the singlet-octet (SO) and the quark flavour (QF) basis; in each scheme two mixing angles are involved [18].

In the SO basis one defines the η and η' -vacuum matrix elements of axial-vector currents:

$$\langle 0 | J_{\mu 5}^a | P(p) \rangle = i f_P^a p_\mu \quad (a = 1, 8) \quad (3.1)$$

where $P = \eta, \eta'$ and $J_{\mu 5}^1, J_{\mu 5}^8$ are $SU(3)_F$ singlet and octet axial-vector quark currents. The four hadronic parameters f_P^a in (3.1) can be written in terms of two angles: θ_1 and θ_8 , and of two decay constants f_1 and f_8 of a pure singlet and octet flavour state. In this scheme, non-vanishing values of the angles $\theta_{1,8}$, as well as the difference $f_8 \neq f_\pi$ are $SU(3)_F$ breaking effects, and the contribution of the axial $U(1)$ anomaly is encoded in the decay constant f_1 .

On the other hand, in the QF basis one defines the axial-vector currents:

$$J_{\mu 5}^q = \frac{1}{\sqrt{2}} (\bar{u} \gamma_\mu \gamma_5 u + \bar{d} \gamma_\mu \gamma_5 d) \quad , \quad J_{\mu 5}^s = \bar{s} \gamma_\mu \gamma_5 s \quad (3.2)$$

and the matrix elements:

$$\langle 0 | J_{\mu 5}^b | P(p) \rangle = i f_P^b p_\mu \quad (b = q, s) \quad (3.3)$$

Also in this case the four parameters f_P^b can be expressed in terms of two angles φ_q and φ_s , and of two decay constants f_q and f_s of states without and with strangeness, respectively [18]. The difference between the mixing angles $\varphi_q - \varphi_s$ is due to OZI-violating effects and is found to be small ($\varphi_q - \varphi_s < 5^\circ$), so that it has been proposed that the approximation of describing the $\eta - \eta'$ mixing in the QF basis and a single mixing angle is convenient [18]. The simplification $\varphi_q \simeq \varphi_s \simeq \varphi$ is supported by a QCD sum rule analysis of the decays $\phi \rightarrow \eta \gamma$ and $\phi \rightarrow \eta' \gamma$ [19]. Here we adopt the quark flavour basis and define:

$$\begin{aligned} |\eta_q\rangle &= \frac{1}{\sqrt{2}} (|\bar{u}u\rangle + |\bar{d}d\rangle) \\ |\eta_s\rangle &= |\bar{s}s\rangle \quad , \end{aligned} \quad (3.4)$$

so that the $\eta - \eta'$ system can be described in terms of the mixing angle $\varphi = \varphi_q = \varphi_s$:

$$\begin{aligned} |\eta\rangle &= \cos \varphi |\eta_q\rangle - \sin \varphi |\eta_s\rangle \\ |\eta'\rangle &= \sin \varphi |\eta_q\rangle + \cos \varphi |\eta_s\rangle \quad . \end{aligned} \quad (3.5)$$

There are several ways to measure φ , namely through the radiative transitions involving a light vector meson V , such as $V \rightarrow \eta^{(\prime)} \gamma$ or $\eta' \rightarrow V \gamma$, or studying the two-photon decays $\eta^{(\prime)} \rightarrow \gamma \gamma$ [20]. A precise result has recently been obtained by the KLOE Collaboration which, measuring the ratio $\frac{\Gamma(\phi \rightarrow \eta' \gamma)}{\Gamma(\phi \rightarrow \eta \gamma)}$ and assuming the flavour basis with a single mixing angle, quotes:

$\varphi = (41.5 \pm 0.3_{stat} \pm 0.7_{syst} \pm 0.6_{th})^\circ$ [21]. We use this value in our study. In the same analysis, KLOE has also allowed for a glue content in η' , modifying the second relation in (3.5):

$$\begin{aligned} |\eta'\rangle &= \cos \varphi_G \sin \varphi |\eta_q\rangle + \cos \varphi_G \cos \varphi |\eta_s\rangle \\ &+ \sin \varphi_G |gluons\rangle \end{aligned} \quad (3.6)$$

where φ_G is the mixing angle for the glue contribution. In this case, considering also the results for the decay widths $\Gamma(\eta' \rightarrow \gamma \gamma)$, $\Gamma(\eta' \rightarrow \rho \gamma)$ and $\Gamma(\eta' \rightarrow \omega \gamma)$, together with $\Gamma(\pi^0 \rightarrow \gamma \gamma)$ and $\Gamma(\omega \rightarrow \pi^0 \gamma)$, KLOE obtains: $\varphi = (39.7 \pm 0.7)^\circ$ and $\cos^2 \varphi_G = 0.86 \pm 0.04$. We comment below on how this measurement affects our predictions.

The mixing parameters are useful to determine the other quantities needed for the description of $B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^- (\nu \bar{\nu})$ transitions, the B_s and $\eta^{(\prime)}$ matrix elements of the operators in (2.1), (2.3). As usual, such matrix elements are parameterized in terms of form factors:

$$\begin{aligned} \langle \eta^{(\prime)}(p') | \bar{s} \gamma_\mu b | B_s(p) \rangle &= (p + p')_\mu F_1^{B_s \rightarrow \eta^{(\prime)}}(q^2) \\ &+ \frac{M_{B_s}^2 - M_{\eta^{(\prime)}}^2}{q^2} q_\mu \left(F_0^{B_s \rightarrow \eta^{(\prime)}}(q^2) - F_1^{B_s \rightarrow \eta^{(\prime)}}(q^2) \right) \end{aligned} \quad (3.7)$$

$$(q = p - p', F_1^{B_s \rightarrow \eta^{(\prime)}}(0) = F_0^{B_s \rightarrow \eta^{(\prime)}}(0)) \text{ and}$$

$$\begin{aligned} \langle \eta^{(\prime)}(p') | \bar{s} i \sigma_{\mu\nu} q^\nu b | B_s(p) \rangle &= \\ \left[(p + p')_\mu q^2 - (M_{B_s}^2 - M_{\eta^{(\prime)}}^2) q_\mu \right] &\frac{F_T^{B_s \rightarrow \eta^{(\prime)}}(q^2)}{M_{B_s} + M_{\eta^{(\prime)}}} \quad . \end{aligned} \quad (3.8)$$

No QCD sum rule or lattice QCD calculations of $B_s \rightarrow \eta^{(\prime)}$ form factors (which probe the $\bar{s}s$ content of η and η') are available, yet. The quark flavour scheme allows to relate the $B_s \rightarrow \eta_s$ form factors to the $B \rightarrow K$ ones, so that: $F^{B_s \rightarrow \eta} = -\sin \varphi F^{B \rightarrow K}$ and $F^{B_s \rightarrow \eta'} = \cos \varphi F^{B \rightarrow K}$ (for a generic form factor F), keeping the physical masses of B_s , η and η' . It is possible to estimate the uncertainty connected to flavour symmetry breaking, considering relations holding in the heavy quark limit and in the chiral limit for heavy-to-light transition form factors [22]. Taking as an example the form factor F_1 , one can write for $q^2 \rightarrow q_{max}^2$:

$$F_1^{H \rightarrow P} = \frac{\hat{F} g \sqrt{M_H}}{2 f_P (E_P + \Delta_H)} \quad (3.9)$$

where H and P are a heavy and a light pseudoscalar meson, respectively, and E_P is the energy of the P meson in the H rest frame. $\hat{F}$ is the heavy meson decay constant in the heavy quark limit, which is, at leading order in the heavy quark mass m_Q , independent of it (modulo logarithms). g describes the effective coupling $H^* H P$, H^* being the vector meson with the same quark content as H ; $\Delta_H = M_{H^*} - M_H$. If the main flavour

breaking effects are in M_H , Δ_H , E_P and f_P , the ratio $F_1^{B_s \rightarrow \eta_s} / F_1^{B \rightarrow K}$, using $f_K = 159.8 \pm 1.4 \pm 0.44$ MeV [23] and $f_s = (1.34 \pm 0.06)f_\pi$ [18], differs from 1 by about 10%, a reasonable estimate of $SU(3)_F$ breaking in the form factors.

In the following, we use two different sets of form factors, both obtained by QCD sum rules [24]. We refer to the first set, computed using short-distance QCD sum rules, as set A [25], and to the second set, computed by light-cone QCD sum rules, as set B [26]. The comparison allows a discussion of the uncertainty related to the hadronic matrix elements. In both sets the error of the form factors is given at $q^2 = 0$ and then extended to the full range of momentum transfer: $4m_\ell^2 \leq q^2 \leq (M_{B_s} - M_{\eta^{(\prime)}})^2$.

We also consider a third determination obtained applying light-cone QCD sum rules within the Soft Collinear Effective Theory (SCET) [27]. It is interesting to consider this framework, since it allows to express the form factors in terms of a single universal function $\xi^{B_a \rightarrow P}$ (a light flavour index) through the relations, holding in the heavy-quark limit and in the large energy limit of the light meson P :

$$\begin{aligned} F_1^{B_a \rightarrow P}(q^2) &= \xi^{B_a \rightarrow P}(n_+ \cdot p') \\ F_0^{B_a \rightarrow P}(q^2) &= \frac{2E'}{M_{B_a}} \xi^{B_a \rightarrow P}(n_+ \cdot p') \\ F_T^{B_a \rightarrow P}(q^2) &= \frac{M_{B_a} + M_P}{M_{B_a}} \xi^{B_a \rightarrow P}(n_+ \cdot p') . \end{aligned} \quad (3.10)$$

In (3.10) n_+ is a light-like four-vector with components $n_+ = (1, 0, 0, -1)$, and p' is the momentum of the P meson, with $E' = p'^0$. At first order in the light pseudoscalar meson mass the following relations hold:

$$\begin{aligned} (n_+ \cdot p') &= 2E' - \frac{M_P^2}{2E'} \\ E' &= \frac{M_{B_a}^2 + M_P^2 - q^2}{2M_{B_a}} . \end{aligned} \quad (3.11)$$

We refer to this third parameterization as to set C. Developing light-cone QCD sum rules in the framework of the SCET, in Ref. [28] the universal function $\xi^{B \rightarrow \pi}$ relative to $B \rightarrow \pi$ form factors was derived. By the same method, a determination can be obtained in the case of kaon, and the resulting function $\xi^{B \rightarrow K}$ can be parameterized as:

$$\xi^{B \rightarrow K}(n_+ \cdot p') = \xi^{B \rightarrow K}(m_B) \left[-a + \frac{b}{(n_+ \cdot p')} + c(n_+ \cdot p') \right] \quad (3.12)$$

with

$$\begin{aligned} \xi^{B \rightarrow K}(m_B) &= 0.335_{-0.094}^{+0.078} \\ a &= 2.418 \quad b = 13.765 \quad c = 0.154 . \end{aligned} \quad (3.13)$$

The $B_s \rightarrow \eta^{(\prime)}$ form factors belonging to set C can be written in terms of the universal function:

$$F_1^{B_s \rightarrow \eta} = -\sin \varphi \xi^{B \rightarrow K}$$

$$\begin{aligned} F_0^{B_s \rightarrow \eta} &= -\sin \varphi \frac{2E'}{M_{B_s}} \xi^{B \rightarrow K} \\ F_T^{B_s \rightarrow \eta} &= -\sin \varphi \frac{M_{B_s} + M_\eta}{M_{B_s}} \xi^{B \rightarrow K} \\ F_1^{B_s \rightarrow \eta'} &= \cos \varphi \xi^{B \rightarrow K} \\ F_0^{B_s \rightarrow \eta'} &= \cos \varphi \frac{2E'}{M_{B_s}} \xi^{B \rightarrow K} \\ F_T^{B_s \rightarrow \eta'} &= \cos \varphi \frac{M_{B_s} + M_{\eta'}}{M_{B_s}} \xi^{B \rightarrow K} , \end{aligned} \quad (3.14)$$

keeping the physical masses of the particles involved in the transition. Although such relations are valid in the heavy quark limit at small values of the momentum transfer q^2 , we extrapolate them in the full kinematical range of $B_s \rightarrow \eta^{(\prime)}$ transitions.

Relating $B_s \rightarrow \eta^{(\prime)}$ form factors to the $B \rightarrow K$ ones in the QF scheme implies the neglect of flavour-singlet contributions, for example the annihilation contribution involving the strange quark spectator in B_s and producing $\eta^{(\prime)}$ through two gluons. This contribution has been considered as possibly responsible of the anomalous pattern of the branching fractions of nonleptonic B and D to η' decays. In particular, one could expect that such a kind of contributions are more important for η' than for η , due to the coupling of η' to gluons driven by the $U(1)_A$ anomaly in QCD (at the origin of the large η' mass compared to the masses of light pseudoscalar mesons).

However, the size of this contribution is not firmly established. Denoting this gluonic contribution as F_G and the quark contribution as F_q , a generic form factor F can be parameterized as:

$$\begin{aligned} F^{B_s \rightarrow \eta} &= -F_q \frac{f_q}{f_K} \sin \varphi + F_G \left(\sqrt{\frac{2}{3}} \frac{f_q}{f_K} \cos \varphi - \frac{1}{\sqrt{3}} \frac{f_s}{f_K} \sin \varphi \right) \\ F^{B_s \rightarrow \eta'} &= F_q \frac{f_q}{f_K} \cos \varphi + F_G \left(\sqrt{\frac{2}{3}} \frac{f_q}{f_K} \sin \varphi + \frac{1}{\sqrt{3}} \frac{f_s}{f_K} \cos \varphi \right) \end{aligned} \quad (3.15)$$

in terms of the decay constants $f_q = (1.02 \pm 0.02)f_\pi$ [18], f_s and f_K ; eqs.(3.15) show that the main effect is for the η' form factors. In case of B meson transitions, various analyses conclude that the singlet term F_G could be sizeable, thus possibly contributing to the processes involving η' [29, 30, 31], however the uncertainties are large. To estimate such contributions the matrix elements:

$$< 0 | A_\mu^a(z) A_\nu^b(0) | \eta^{(\prime)}(p) > \quad (3.16)$$

are needed (A_μ^a is the gluon field, and a gauge factor has been dropped). These matrix elements can be written in terms of distribution amplitudes (DA) expanded in Gegenbauer polynomials. However, already the first coefficient (denoted by B_2) of the expansion of the twist-two distribution amplitude ϕ^G (assumed to be the same for η and η') is uncertain: for example, $B_2(1.4 \text{ GeV}) = 4.6 \pm 2.5$ results from a combined analysis of the $\eta' \eta \gamma$ form factor and of the inclusive decay $Y(1S) \rightarrow \eta' X$ [32]. In correspondence to this value a contribution of about 5% is

estimated for the form factor $F_1^{B \rightarrow \eta'}(0)$ [31], and only if B_2 is varied in a wider range ($B_2 = 0 \pm 20$) up to $\mathcal{O}(20\%)$ corrections are found in case of η' , while the corrections remain small in the case of η meson. In the framework of SCET these contributions are found to be consistent with zero [33]. A constraint of the singlet contribution using the widths of the semileptonic decays $B^+ \rightarrow \eta \ell^+ \nu_\ell$ and $B^+ \rightarrow \eta' \ell^+ \nu_\ell$ is hampered by the present experimental errors.

One can investigate the presence of gluonic contributions to the form factors governing the semileptonic modes $D_s \rightarrow \eta \ell^+ \nu_\ell$ and $D_s \rightarrow \eta' \ell^+ \nu_\ell$. The measurements $BR(D_s \rightarrow \eta \ell^+ \nu_\ell) = (3.2 \pm 0.5) \times 10^{-2}$ and $BR(D_s \rightarrow \eta' \ell^+ \nu_\ell) = (1.12 \pm 0.35) \times 10^{-2}$ [23] can be combined in the ratio: $R_{D_s} = \frac{BR(D_s \rightarrow \eta' \ell^+ \nu_\ell)}{BR(D_s \rightarrow \eta \ell^+ \nu_\ell)} = 0.35 \pm 0.12$. This value for R_{D_s} is reproduced by central values of the mixing angle: $\varphi = 41.5^\circ$ and of the ratio of the two pieces

contributing to $F_1^{D_s \rightarrow \eta^{(\prime)}}(0)$ written in (3.15): $\frac{F_G(0)}{F_q(0)} = 0$, thus pointing to small values of the singlet terms.

A possibility for a phenomenological analysis of $B_s \rightarrow \eta, \eta'$ transitions could consist in considering the effects of an arbitrarily added flavour singlet contribution to the form factors [34]. We prefer to provide predictions which do not include additional terms in the form factors. The predictions are accurate in case of η ; deviations in the modes with η' would first prompt investigations focused on the singlet contributions.

IV. $B_s \rightarrow \eta \ell^+ \ell^-$ AND $B_s \rightarrow \eta' \ell^+ \ell^-$

The observables in these modes are the differential and total decay rates. The expression for the differential width:

$$\frac{d\Gamma(B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-)}{dq^2} = \frac{G_F^2 |V_{tb} V_{ts}^*|^2 \alpha^2}{2^9 \pi^5} \frac{\lambda^{1/2}(M_{B_s}^2, M_{\eta^{(\prime)}}^2, q^2)}{M_{B_s}^3} \sqrt{1 - \frac{4m_\ell^2}{q^2}} \frac{1}{3q^2} p(q^2) \quad (4.1)$$

($\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2xz - 2yz$) involves the function

$$p(q^2) = 6m_\ell^2(M_{B_s}^2 - M_{\eta^{(\prime)}}^2)^2 |\tilde{b}(q^2)|^2 + \lambda(M_{B_s}^2, M_{\eta^{(\prime)}}^2, q^2) [(2m_\ell^2 + q^2) |\tilde{c}(q^2)|^2 - (4m_\ell^2 - q^2) |\tilde{a}(q^2)|^2] \quad (4.2)$$

where $\tilde{a}$, $\tilde{b}$ and $\tilde{c}$ are combinations of form factors and Wilson coefficients:

$$\begin{aligned} \tilde{a}(q^2) &= C_{10} F_1^{B_s \rightarrow \eta^{(\prime)}}(q^2) \\ \tilde{b}(q^2) &= C_{10} F_0^{B_s \rightarrow \eta^{(\prime)}}(q^2) \\ \tilde{c}(q^2) &= C_9 F_1^{B_s \rightarrow \eta^{(\prime)}}(q^2) \\ &\quad - 2(m_b + m_s) C_7^{eff} \frac{F_T^{B_s \rightarrow \eta^{(\prime)}}(q^2)}{M_{B_s} + M_{\eta^{(\prime)}}} \end{aligned} \quad (4.3)$$

In the expression of $\tilde{c}$ in (4.3) we introduced the coefficient C_7^{eff} , which is a renormalization scheme independent combination of C_7, C_8 and C_2 , given by a formula that can be found, e.g., in [9]. Using: $m_b = 4.8 \pm 0.2$ GeV, $m_s = 104_{-34}^{+26}$ MeV, together with the values: $M_{B_s^0} = 5366.3 \pm 0.6$ MeV, $\tau_{B_s^0} = (1.470_{-0.027}^{+0.026}) \times 10^{-12}$ s, $V_{tb} = 0.999$ and $V_{ts} = (38.7 \pm 2.3) \times 10^{-3}$ quoted by the Particle Data Group [23], we obtain the following branching ratios in the SM, depending on the set of form factors:

$$BR(B_s \rightarrow \eta \ell^+ \ell^-) = \begin{cases} (1.2 \pm 0.3) \times 10^{-7} & \text{set A} \\ (2.6 \pm 0.7) \times 10^{-7} & \text{set B} \\ (3.4 \pm 1.8) \times 10^{-7} & \text{set C} \end{cases} \quad (4.4)$$

$$BR(B_s \rightarrow \eta' \ell^+ \ell^-) = \begin{cases} (1.1 \pm 0.3) \times 10^{-7} & \text{set A} \\ (2.2 \pm 0.6) \times 10^{-7} & \text{set B} \\ (2.8 \pm 1.5) \times 10^{-7} & \text{set C} \end{cases} \quad (4.5)$$

for $\ell = e, \mu$, and

$$BR(B_s \rightarrow \eta \tau^+ \tau^-) = \begin{cases} (3 \pm 0.5) \times 10^{-8} & \text{set A} \\ (8 \pm 1.5) \times 10^{-8} & \text{set B} \\ (10 \pm 5.5) \times 10^{-8} & \text{set C} \end{cases} \quad (4.6)$$

$$BR(B_s \rightarrow \eta' \tau^+ \tau^-) = \begin{cases} (1.55 \pm 0.3) \times 10^{-8} & \text{set A} \\ (3.85 \pm 0.75) \times 10^{-8} & \text{set B} \\ (4.7 \pm 2.5) \times 10^{-8} & \text{set C} \end{cases} \quad (4.7)$$

The results obtained using the form factor $\xi^{B \rightarrow K}$ in the SCET framework are affected by the largest uncertainty and encompass the predictions based on the set A and set B. We consider these results as the most conservative determinations of the decay rates, as well as for the other observables.

The results from set A and B are compatible within the errors, with lower central values for set A. The uncertainties, estimated at the level of 30% in the rates for each set of form factor, become higher when the central values are compared: the difference provides us with hints on the level of improvement in the determination of the form factors needed at present. Such an improvement




FIG. 1: Differential branching fraction $dBR(B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-)/dq^2$ ($\ell = e, \mu$) as a function of the momentum transfer q^2 (upper plots), and branching fraction $BR(B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-)$ versus the compactification scale R^{-1} (lower plots) obtained using set A and B of form factors. In each upper plot the lowest (red) region is obtained in SM ($1/R \rightarrow \infty$), the intermediate (blue) region for the compactification scale $1/R = 500 \text{ GeV}$ and the uppermost (green) region for $1/R = 200 \text{ GeV}$.

also concerns other determinations of the form factors [35].

As a further remark, we notice that the KLOE fit of the $\eta - \eta'$ mixing eq.(3.6) corresponds to using an effective mixing angle $\varphi^{eff} = (44.5 \pm 1.5)^\circ$ in the expressions of the $B_s \rightarrow \eta'$ form factors, and to a 10% decrease of the corresponding decay rates.

It is worth comparing the predictions (4.4)-(4.5) with the results quoted by the Heavy Flavour Averaging Group (HFAG) for the analogous B decay modes [36]: $BR(B^0 \rightarrow K^0 \ell^+ \ell^-) = (3.2^{+0.8}_{-0.6}) \times 10^{-7}$ and $BR(B^+ \rightarrow K^+ \ell^+ \ell^-) = (4.9 \pm 0.5) \times 10^{-7}$ for $\ell = e, \mu$. On the other hand, in the approach based on the QF mixing scheme, a relation connecting the decay width of $B_s \rightarrow \eta \ell^+ \ell^-$ to that of $B_s \rightarrow K^0 \ell^+ \ell^-$ induced by the transition $b \rightarrow d$:

$$\frac{BR(B_s \rightarrow K^0 \ell^+ \ell^-)}{BR(B_s \rightarrow \eta \ell^+ \ell^-)} = \frac{1}{(\sin \varphi)^2} \left| \frac{V_{td}}{V_{ts}} \right|^2 (1 + \Delta_{SU(3)} + \Delta'), \quad (4.8)$$

involves an $SU(3)$ correction term $\Delta_{SU(3)}$ (presumably small) and a term Δ' coming from the additional contribution proportional to $V_{ub}V_{ud}^*$, the correspondent of which has been neglected in the effective Hamiltonian (2.1) in case of $b \rightarrow s$ modes. Such a kind of relations is of great interest, both from the theoretical and the experimental viewpoint, and deserves a dedicated analysis.

In the ACD model the Wilson coefficients C_7^{eff} , C_9 and C_{10} depend on the compactification scale $1/R$, hence decay rates and distributions vary with this parameter. C_7^{eff} depends on $1/R$ through two functions $D'(x_t, \frac{1}{R})$ and $E'(x_t, \frac{1}{R})$ representing the contribution of γ -penguins and chromomagnetic penguins, respectively. The contribution of Z^0 -penguins enters in the functions $Y(x_t, \frac{1}{R})$, $Z(x_t, \frac{1}{R})$ and $X(x_t, \frac{1}{R})$: Y and Z determine the

coefficient C_9 , while C_{10} depends only on Y . As for the function X , it enters in the ACD expression of the coefficient C_L relevant for the modes with neutrinos discussed in the next Section.

These functions obey the general representation (2.5), and can be found, e.g., in [8, 9]. For values of the compactification scale $1/R$ of a few hundreds of GeV the coefficient C_7^{eff} is suppressed in the ACD model with respect to the SM value, while C_{10} is enhanced and C_9 turns out to be almost unaffected. For example, at $\frac{1}{R} = 300 \text{ GeV}$ one has: $\frac{C_7^{eff,ACD}}{C_7^{eff,SM}} \simeq 0.82$, $\frac{C_9^{ACD}}{C_9^{SM}} \simeq 1.01$, $\frac{C_{10}^{ACD}}{C_{10}^{SM}} \simeq 1.16$. As a consequence, for the decay into η and $m_\ell = 0$ we find for the function $p(q^2)$ in (4.2) at $q^2 = q_{max}^2$ and $1/R = 300 \text{ GeV}$: $\frac{p(q_{max}^2)|_{1/R=300 \text{ GeV}}}{p(q_{max}^2)|_{SM}} = 1.23 - 1.42$, depending on the set of form factors.

The dependence of the rates on $1/R$ is obscured by the uncertainty of the hadronic form factors, as shown in Fig. 1, at values of this scale excluded by the analyses of the electroweak constraints: $1/R \geq 250 \text{ GeV}$. Sensitivity of the decay rate to $1/R$ could be achieved by a reduction in the hadronic uncertainty of about a factor of two: in this case, the measurement of the rates of $B_s \rightarrow \eta^{(\prime)} \ell^+ \ell^-$ would allow to exclude the $1/R$ region below $350 - 400 \text{ GeV}$, keeping the same uncertainties quoted for the other input parameters. Concerning the differential distributions $\frac{d\Gamma}{dq^2}(B_s \rightarrow \eta \ell^+ \ell^-)$ and $\frac{d\Gamma}{dq^2}(B_s \rightarrow \eta' \ell^+ \ell^-)$ also depicted in Fig.1, the enhancement at small values of dilepton invariant mass obtained for decreasing compactification scales $1/R$ could be observed after the same improvement of the accuracy of the form factors.

A possibility to reduce the effects of the hadronic uncertainties consists in comparing the decay distributions of $B_s \rightarrow \eta \ell^+ \ell^-$ and $D_s \rightarrow \eta \ell^+ \nu_\ell$ ($\ell = e, \mu$) at large values of momentum transfer $q^2 \simeq q_{max}^2$, i.e. close to the end-point. In this range the ratio:

$$R(y) = \frac{d\Gamma(B_s \rightarrow \eta \ell^+ \ell^-)/dy}{d\Gamma(D_s \rightarrow \eta \ell^+ \nu_\ell)/dy} \quad (4.9)$$

obtained evaluating the distributions at the same value of $y = \frac{E}{M_\eta}$ can be written, up to corrections $\mathcal{O}(\frac{1}{m_c} - \frac{1}{m_b})$, as:

$$R(y) = \left| \frac{V_{tb} V_{ts}^*}{V_{cs}} \right|^2 \frac{\alpha^2}{8\pi^2} \left(\frac{M_{B_s}}{M_{D_s}} \right)^2 \left(|C_{10}|^2 + \left| C_9 - \frac{2(m_b + m_s)}{M_{B_s} + M_\eta} c_7^{eff} \frac{F_T^{B_s \rightarrow \eta}(y)}{F_1^{B_s \rightarrow \eta}(y)} \right|^2 \right) \quad (4.10)$$

thus providing an access to the Wilson coefficients $C_{7,9}$ and C_{10} , since the ratio of $\frac{F_T^{B_s \rightarrow \eta}(y)}{F_1^{B_s \rightarrow \eta}(y)}$ is practically independent of y and of the set of form factors. To further reduce the impact of the hadronic corrections, the double ratio could be considered:

$$\tilde{R}(y) = \left[\frac{d\Gamma(B_s \rightarrow \eta \ell^+ \ell^-)/dy}{d\Gamma(D_s \rightarrow \eta \ell^+ \nu_\ell)/dy} \right] / \left[\frac{d\Gamma(B_s \rightarrow K \ell^+ \nu)/dy}{d\Gamma(D_s \rightarrow K \ell^+ \nu_\ell)/dy} \right] \quad (4.11)$$

which is given by

$$\tilde{R}(y) = \left| \frac{V_{tb} V_{ts}^*}{V_{cs}} \right|^2 \left| \frac{V_{cd}}{V_{ub}} \right|^2 \frac{\alpha^2}{8\pi^2} \left(\frac{M_{B_s}}{M_{D_s}} \right)^2 \left(|C_{10}|^2 + \left| C_9 - \frac{2(m_b + m_s)}{M_{B_s} + M_\eta} c_7^{eff} \frac{F_T^{B_s \rightarrow \eta}(y)}{F_1^{B_s \rightarrow \eta}(y)} \right|^2 \right) \quad (4.12)$$

up to corrections $\mathcal{O}(m_s(\frac{1}{m_c} - \frac{1}{m_b}))$ [37]. However, the double ratio involves many CKM suppressed transitions, and its measurement is challenging.

V. $B_s \rightarrow \eta \nu \bar{\nu}$ AND $B_s \rightarrow \eta' \nu \bar{\nu}$

The exclusive decays with two neutrinos in the final state, which from the theoretical side are among the cleanest FCNC processes, require only one hadronic form factor. The observables are the decay distributions and the total decay widths. For the former, it is convenient to use the variable $x = \frac{E_{miss}}{M_{B_s}}$, with E_{miss} the energy of the neutrino pair in the B_s rest frame (missing energy), with $0.5 \leq x \leq 1 - \frac{M_{\eta^{(\prime)}}}{M_{B_s}}$. In terms of the variable x the

differential decay rate $\frac{d\Gamma}{dx}$ reads:

$$\frac{d\Gamma(B_s \rightarrow \eta^{(\prime)} \nu \bar{\nu})}{dx} = 3 \frac{|C_L|^2 |F_1^{B_s \rightarrow \eta^{(\prime)}}(q^2)|^2}{48\pi^3 M_{B_s}} \sqrt{\lambda^3(q^2, M_{B_s}^2, M_{\eta^{(\prime)}}^2)}, \quad (5.1)$$

with the invariant mass q^2 of the neutrino pair expressed in terms of x : $q^2 = M_{B_s}^2(2x-1) + M_{\eta^{(\prime)}}^2$. C_L is the Wilson coefficient in (2.3) and the factor of 3 corresponds to the sum over the three neutrino flavours.

The differential branching fraction is plotted in Fig. 2 in the case of η and η' , respectively, for the sets A and B of form factors. In the Standard Model one predicts:

$$BR(B_s \rightarrow \eta \nu \bar{\nu}) = \begin{cases} (0.95 \pm 0.2) \times 10^{-6} & \text{set A} \\ (2.2 \pm 0.7) \times 10^{-6} & \text{set B} \\ (2.9 \pm 1.5) \times 10^{-6} & \text{set C} \end{cases} \quad (5.2)$$

$$BR(B_s \rightarrow \eta' \nu \bar{\nu}) = \begin{cases} (0.9 \pm 0.2) \times 10^{-6} & \text{set A} \\ (1.9 \pm 0.5) \times 10^{-6} & \text{set B} \\ (2.4 \pm 1.3) \times 10^{-6} & \text{set C} \end{cases} \quad (5.3)$$

These predictions can be compared to the present upper bounds for the corresponding $B \rightarrow K \nu \bar{\nu}$ modes, quoted by HFAG: $BR(B^0 \rightarrow K^0 \nu \bar{\nu}) < 160 \times 10^{-6}$ and $BR(B^+ \rightarrow K^+ \nu \bar{\nu}) < 14 \times 10^{-6}$ [36], which are compatible with the SM expectations [38].

In the ACD model, the coefficient is slightly enhanced: for example $\frac{C_L^{ACD}}{C_L^{SM}} \simeq 1.10$ at $1/R = 300$ GeV. However, the $1/R$ dependence of the branching fractions is obscured by the hadronic uncertainty, as shown in Fig. 2. The distribution in missing energy shows an enhancement at small values of x which becomes sizeable at the lowest values of the $1/R$ range, $1/R \sim 250$ GeV. The presence of two neutrinos in the final state makes the measurements of these modes a challenge that can be faced at an e^+e^- superB factory operating at the $\Upsilon(5S)$ peak.

VI. THE bs UT TRIANGLE IN ACD

To complete our discussion about B_s physics in the ACD model, we turn to the bs unitarity triangle (UT) and, in particular, to the weak phase β_s defined as: $\beta_s = \text{Arg} \left[-\frac{V_{ts} V_{tb}^*}{V_{cs} V_{cb}^*} \right]$. Being ACD a Minimal Flavour Violation model, the CKM matrix has the same structure as in the SM: it is constrained to be unitary and is described in terms of four parameters, one of which is a complex phase: ρ , η , A and λ in the Wolfenstein parameterization [39]. At $\mathcal{O}(\lambda^3)$ only V_{ub} and V_{td} have a complex phase, while at $\mathcal{O}(\lambda^4)$ also V_{ts} is complex and its argument identifies β_s [40].




FIG. 2: Differential decay rates $dBR(B_s \rightarrow \eta^{(i)} \nu \bar{\nu})/dx$ as a function of $x = E_{miss}/M_{B_s}$ (upper plots), and branching fractions $BR(B_s \rightarrow \eta^{(i)} \nu \bar{\nu})$ versus R^{-1} (lower plots) obtained using set A and B (right) of form factors. In the upper plots the lowest region is obtained in SM; the intermediate (blue) one for $1/R = 500$ GeV, the uppermost (green) one for $1/R = 200$ GeV.

As it is well known, the various unitarity conditions are represented as triangles in the complex plane. Even though the CKM matrix has the same structure in ACD and SM, and, in particular, its phase is in both cases the only source of CP violation, it may happen that the various CKM elements differ in the two models and therefore the UT triangles do not coincide. Deviations could be expected for elements extracted from loop-induced processes, where the tower of KK modes may play a role; for elements obtained from tree level decays no difference is expected, as for $|V_{us}|$, $|V_{ub}|$ and $|V_{cb}|$. For the db triangle defined by the relation: $V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$, only the side $R_t = \frac{|V_{td}V_{tb}^*|}{|V_{cd}V_{cb}^*|}$ depends on R^{-1} through the dependence of $|V_{td}|$ coming from the KK contribution to the $B_d^0 - \bar{B}_d^0$ mixing [8]. Using such a result, together with the measured value of $\frac{|V_{ub}|}{|V_{cb}|}$, the variation

of $\bar{\rho} = \left(1 - \frac{\lambda^2}{2}\right)\rho$, $\bar{\eta} = \left(1 - \frac{\lambda^2}{2}\right)\eta$, and of γ (the phase of V_{ub}^*) has been obtained [8], finding that the db triangle in the ACD model could be slightly different than in the SM.

The bs triangle stems from the unitarity condition involving CKM elements relevant for B_s decays, and is defined by the relation:

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0. \quad (6.1)$$

Analogously to V_{td} , V_{ts} inherits the R^{-1} dependence from $B_s^0 - \bar{B}_s^0$ mixing [8]: $\Delta M_s \propto S(x_t, 1/R)|V_{ts}|^2$, where the proportionality factor does not depend on R^{-1} and is common to ACD and SM. The function $S(x_t, 1/R)$ takes into account the SM contribution, $S_0(x_t)$, and the contribution of the KK modes: $S(x_t, 1/R) = S_0(x_t) + \sum_n S_n(x_t, 1/R)$, with:

$$\begin{aligned} \sum_n S_n(x_t, x_n) = & -\frac{x_t(5 - 10x_t + x_t^2)}{8(x_t - 1)^2} + \frac{\pi M_W R}{2} \left\{ -\frac{x_t(3x_t - 1)}{2(x_t^2 - 1)^3} J(R, -1/2) + \frac{(1 - 3x_t + 7x_t^2 - x_t^3)}{2(x_t - 1)^3} J(R, 1/2) \right. \\ & \left. - \frac{(-3 + 6x_t - x_t^3)}{2x_t(x_t - 1)^3} J(R, 3/2) + \frac{1}{2x_t} \int_0^1 dy [x_t + 3(1 + x_t)y] \sqrt{y} \operatorname{ctanh}(\pi M_W R \sqrt{y}) \right\} \end{aligned} \quad (6.2)$$

and

As a consequence one finds:

$$|V_{ts}|_{ACD} = |V_{ts}|_{SM} \sqrt{\frac{S_0(x_t)}{S(x_t, 1/R)}}. \quad (6.4)$$

$J(R, \alpha) = \int_0^1 dy y^\alpha [\operatorname{ctanh}(\pi M_W R \sqrt{y}) - x_t^{1+\alpha} \operatorname{ctanh}(\pi m_t R \sqrt{y})]$. In Fig.3 we show the dependence of $|V_{ts}|_{ACD}$ on R^{-1} using the quoted value of $|V_{ts}|_{SM}$. Since at $O(\lambda^4)$




FIG. 3: CKM element V_{ts} (left) and phase β_s (rad) (right) versus R^{-1} .

$V_{ts} = -A\lambda^2 + \frac{A\lambda^4}{2}[1 - 2\rho - 2i\eta]$, it turns out that $\beta_s = \arctan \left[\frac{2\lambda^2\eta}{2 - \lambda^2(1 - 2\rho)} \right]$. Using the dependence of ρ and η (or $\bar{\rho}$ and $\bar{\eta}$), also the β_s dependence on R^{-1} can be obtained; it is depicted in Fig. 3. The SM result for β_s is small: $\beta_s \simeq 0.019$ rad, and the dependence on the compactification scale R^{-1} further reduces this value. As mentioned in the Introduction, preliminary Tevatron results point towards larger values of β_s , so that our analysis supports the conclusion that a confirmation of the measurement of a large phase in the B_s mixing would point towards new Physics models different from MFV scenarios.

In Fig. 4 the rescaled bs UT triangle in the ACD model is displayed at the representative compactification scale $R^{-1} = 300$ GeV.




FIG. 4: Rescaled bs UT triangle in the SM (continuous line) and the ACD model at $R^{-1} = 300$ GeV (dashed line).

VII. CONCLUSIONS

Together with other rare B_s decays, the loop-induced transitions $B_s \rightarrow \eta^{(\prime)}\ell^+\ell^-$ and $B_s \rightarrow \eta^{(\prime)}\nu\bar{\nu}$ must be included in the Physics programmes of experiments aimed at performing tests of the Standard Model and at searching signals of new Physics like those related to extra dimensions. Within SM, using the flavour scheme for the $\eta - \eta'$ mixing and the angle determined by the KLOE Collaboration from radiative ϕ decays, we have predicted decay widths and distributions of these rare B_s channels.

The results for the branching fractions of modes with two charged leptons are $\mathcal{O}(10^{-7})$ for $\ell = e, \mu$ and $\mathcal{O}(10^{-8})$ for $\ell = \tau$, suggesting that they are within the reach of facilities like a SuperB factory. In the case of the final states with two neutrinos, the branching ratios are of $\mathcal{O}(10^{-6})$. The uncertainty in the numerical results is dominated by the error of the hadronic form factors: in order to be sensitive to compactification scales $1/R \sim 350 - 400$ GeV in the ACD model the error in the form factors should be reduced by about a factor of two.

A remark concerns the issue of the $\eta - \eta'$ mixing. In addition to modes with light mesons, processes involving heavy mesons provide information on this mixing. Examples are the J/ψ radiative decays: $J/\psi \rightarrow \eta'\gamma$ and $J/\psi \rightarrow \eta\gamma$, which are sensitive to the glue content of η and η' [41]. Although experimentally challenging and considered mainly for different purposes, also the ratio $\frac{BR(B_s \rightarrow \eta'\ell^+\ell^-)}{BR(B_s \rightarrow \eta\ell^+\ell^-)}$ represents a possibility to determine φ and to look at the gluonic content of η' . A discrepancy in the comparison of the determination of the same quantity from other channels, e.g. from the ratio $\frac{BR(\phi \rightarrow \eta'\gamma)}{BR(\phi \rightarrow \eta\gamma)}$, would signal peculiar effects in $B_s \rightarrow \eta^{(\prime)}$ modes.

Acknowledgments

FDF thanks Thorsten Feldmann and Tobias Hurth for collaboration on light-cone QCD sum rules in the SCET framework. This work was supported in part by the EU contract No. MRTN-CT-2006-035482, "FLAVIANet".

-
- [1] P. Ball *et al.*, “ B Decays at the LHC,” arXiv:hep-ph/0003238; K. Anikeev *et al.*, “ B physics at the Tevatron: Run II and beyond,” arXiv:hep-ph/0201071; M. Bona *et al.*, “SuperB: A High-Luminosity Asymmetric e^+e^- Super Flavor Factory. Conceptual Design Report,” arXiv:0709.0451 [hep-ex]; M. Artuso *et al.*, “ B , D and K Decays”, Eur. Phys. J. C **57**, 309 (2008).
- [2] V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **98**, 121801 (2007); Phys. Rev. D **76**, 057101 (2007); T. Aaltonen *et al.* [CDF collaboration], Phys. Rev. Lett. **100**, 121803 (2008); Phys. Rev. Lett. **100**, 161802 (2008).
- [3] A. Drutskoy, arXiv:0905.2959 [hep-ex].
- [4] T. Appelquist, H. C. Cheng and B. A. Dobrescu, Phys. Rev. D **64**, 035002 (2001).
- [5] T. Appelquist and H. U. Yee, Phys. Rev. D **67**, 055002 (2003).
- [6] H. C. Cheng, K. T. Matchev and M. Schmaltz, Phys. Rev. D **66**, 036005 (2002), Phys. Rev. D **66**, 056006 (2002); G. Servant and T. M. P. Tait, Nucl. Phys. B **650**, 391 (2003).
- [7] For a review see: D. Hooper and S. Profumo, Phys. Rept. **453**, 29 (2007).
- [8] A. J. Buras, M. Spranger and A. Weiler, Nucl. Phys. B **660**, 225 (2003); A. J. Buras, A. Poschenrieder, M. Spranger and A. Weiler, Nucl. Phys. B **678**, 455 (2004).
- [9] P. Colangelo, F. De Fazio, R. Ferrandes and T. N. Pham, Phys. Rev. D **73**, 115006 (2006); Phys. Rev. D **74**, 115006 (2006); Phys. Rev. D **77**, 055019 (2008).
- [10] U. Haisch and A. Weiler, Phys. Rev. D **76**, 034014 (2007).
- [11] R. Mohanta and A. K. Giri, Phys. Rev. D **75**, 035008 (2007); T. M. Aliev and M. Savci, Eur. Phys. J. C **50**, 91 (2007); T. M. Aliev, M. Savci and B. B. Sirvanli, Eur. Phys. J. C **52**, 375 (2007); I. Ahmed, M. A. Paracha and M. J. Aslam, Eur. Phys. J. C **54**, 591 (2008); A. Sadique, M. J. Aslam and C. D. Lu, Eur. Phys. J. C **56**, 267 (2008); V. Bashiry and K. Zeynali, Phys. Rev. D **79**, 033006 (2009); Y. M. Wang, M. J. Aslam and C. D. Lu, Eur. Phys. J. C **59**, 847 (2009).
- [12] G. Devidze, A. Liparteliani and U. G. Meissner, Phys. Lett. B **634**, 59 (2006); I. I. Bigi, G. G. Devidze, A. G. Liparteliani and U. G. Meissner, Phys. Rev. D **78**, 097501 (2008).
- [13] C. Bobeth, M. Misiak and J. Urban, Nucl. Phys. B **574**, 291 (2000); H. H. Asatrian, H. M. Asatrian, C. Greub and M. Walker, Phys. Lett. B **507**, 162 (2001); Phys. Rev. D **65**, 074004 (2002); Phys. Rev. D **66**, 034009 (2002); H. M. Asatrian, K. Bieri, C. Greub and A. Hovhannisyan, Phys. Rev. D **66**, 094013 (2002); A. Ghinculov, T. Hurth, G. Isidori and Y. P. Yao, Nucl. Phys. B **648**, 254 (2003); A. Ghinculov, T. Hurth, G. Isidori and Y. P. Yao, Nucl. Phys. B **685**, 351 (2004); C. Bobeth, P. Gambino, M. Gorbahn and U. Haisch, JHEP **0404**, 071 (2004).
- [14] T. Inami and C. S. Lim, Prog. Theor. Phys. **65**, 297 (1981) [Erratum-ibid. **65**, 1772 (1981)].
- [15] G. Buchalla and A. J. Buras, Nucl. Phys. B **400**, 225 (1993); G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).
- [16] M. Misiak and J. Urban, Phys. Lett. B **451**, 161 (1999).
- [17] G. Buchalla and A. J. Buras, Nucl. Phys. B **548**, 309 (1999).
- [18] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D **58**, 114006 (1998); Phys. Lett. B **449**, 339 (1999); T. Feldmann, Int. J. Mod. Phys. A **15**, 159 (2000).
- [19] F. De Fazio and M. R. Pennington, JHEP **0007**, 051 (2000).
- [20] R. Escribano, Acta Phys. Polon. Supp. **2**, 71 (2009).
- [21] F. Ambrosino *et al.* [KLOE Collaboration], Phys. Lett. B **648**, 267 (2007).
- [22] M. B. Wise, Phys. Rev. D **45**, 2188 (1992).
- [23] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [24] For a review see: P. Colangelo and A. Khodjamirian, arXiv:hep-ph/0010175.
- [25] P. Colangelo, F. De Fazio, P. Santorelli and E. Scrimieri, Phys. Rev. D **53**, 3672 (1996) [Erratum-ibid. D **57**, 3186 (1998)].
- [26] P. Ball and R. Zwicky, Phys. Rev. D **71**, 014015 (2005).
- [27] C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D **63**, 114020 (2001); C. W. Bauer and I. W. Stewart, Phys. Lett. B **516**, 134 (2001); M. Beneke, A. P. Chapovsky, M. Diehl and T. Feldmann, Nucl. Phys. B **643**, 431(2002); M. Beneke and T. Feldmann, Phys. Lett. B **553**, 267 (2003).
- [28] F. De Fazio, T. Feldmann and T. Hurth, Nucl. Phys. B **733**, 1 (2006) [Erratum-ibid. B **800**, 405 (2008)]; JHEP **0802**, 031 (2008).
- [29] M. Beneke and M. Neubert, Nucl. Phys. B **651**, 225 (2003).
- [30] Y. Y. Charng, T. Kurimoto and H. n. Li, Phys. Rev. D **74**, 074024 (2006) [Phys. Rev. D **78**, 059901 (2008)].
- [31] P. Ball and G. W. Jones, JHEP **0708**, 025 (2007).
- [32] A. Ali and A. Y. Parkhomenko, Eur. Phys. J. C **30**, 183 (2003).
- [33] A. R. Williamson and J. Zupan, Phys. Rev. D **74**, 014003 (2006) [Erratum-ibid. D **74**, 03901 (2006)].
- [34] C. H. Chen and C. Q. Geng, Phys. Lett. B **645**, 197 (2007).
- [35] P. Z. Skands, JHEP **0101**, 008 (2001); C. Q. Geng and C. C. Liu, J. Phys. G **29**, 1103 (2003).
- [36] E. Barberio *et al.* [Heavy Flavor Averaging Group], arXiv:0808.1297 [hep-ex].
- [37] B. Grinstein and D. Pirjol, Phys. Lett. B **549**, 314 (2002).
- [38] P. Colangelo, F. De Fazio, P. Santorelli and E. Scrimieri, Phys. Lett. B **395**, 339 (1997); G. Buchalla, G. Hiller and G. Isidori, Phys. Rev. D **63**, 014015 (2001); W. Altmannshofer, A. J. Buras, D. M. Straub and M. Wick, JHEP **0904**, 022 (2009).
- [39] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).
- [40] A. J. Buras, M. E. Lautenbacher and G. Ostermaier, Phys. Rev. D **50**, 3433 (1994).
- [41] P. Ball, J. M. Frere and M. Tytgat, Phys. Lett. B **365**, 367 (1996).
- [42] The ACD Lagrangian may include, in addition to $5-d$ bulk terms, boundary terms introducing additional parameters in the theory. Such terms are renormalized by bulk interactions, but they are volume suppressed. A simplifying assumption is that they vanish at the cutoff scale, so that the only new parameter with respect to SM is the radius R of the compactified extra dimension.